arXiv:0710.1583v1 [math.NT] 8 Oct 2007

# MANIN'S CONJECTURE FOR A QUINTIC DEL PEZZO SURFACE WITH $\mathbf{A}_2$ SINGULARITY

ULRICH DERENTHAL

Abstract. Manin's conjecture is proved for a split del Pezzo surface of degree 5 with a singularity of type $\mathbf{A}_2$.

## Contents

1. Introduction 1
2. A universal torsor 2
3. Estimating $N_{b_2}(B;A)$ 4
4. Real-valued functions 5
5. Estimating $N_a(B)$ and $N_{b_1}(B;A)$ – first step 6
6. Estimating $N_a(B)$ – second step 7
7. Estimating $N_{b_1}(B;A)$ – second step 11
8. The final step 13

References 16

## 1. Introduction

Let $S \subset \mathbb{P}^5$ be the del Pezzo surface of degree 5 defined by

$$\begin{aligned} & x_0x_2 - x_1x_5 = x_0x_2 - x_3x_4 = x_0x_3 + x_1^2 + x_1x_4 \\ & = x_0x_5 + x_1x_4 + x_4^2 = x_3x_5 + x_1x_2 + x_2x_4 = 0. \end{aligned} \tag{1.1}$$

It contains a unique singularity of type $\mathbf{A}_2$ and four lines, all of them defined over $\mathbb{Q}$. Let $U \subset S$ be the complement of these lines.

We define the height of any rational point $\mathbf{x} \in S(\mathbb{Q})$ that is represented by integral and relatively coprime coordinates $(x_0, \dots, x_5)$ as

$$H(\mathbf{x}) := \max\{|x_0|, \dots, |x_5|\}.$$

For any $B \geqslant 1$, let

$$N_{U,H}(B) := \#\{\mathbf{x} \in U(\mathbb{Q}) \mid H(\mathbf{x}) \leqslant B\}$$

be the number of rational points in $U$ whose height is at most $B$.

We prove the following result:

**Theorem.** *We have*

$$N_{U,H}(B) = c_{S,H}B(\log B)^4 + O(B(\log B)^{4-1/5}),$$

---

2000 *Mathematics Subject Classification.* Primary 11G35; Secondary 14G05.

*where*

$$c_{S,H} = \frac{1}{864} \cdot \omega_\infty \cdot \prod_p \left(1 - \frac{1}{p}\right)^5 \left(1 + \frac{5}{p} + \frac{1}{p^2}\right)$$

*and*

$$\omega_\infty = \int_{|t_5|,|t_1|,|t_1t_5^2t_6^2+t_1^2t_6|,|t_1t_5t_6|,|t_5^2t_6+t_1|,|t_5^3t_6^2+t_1t_5t_6|\leqslant 1,\ t_5>0} \mathrm{d}t_1 \, \mathrm{d}t_5 \, \mathrm{d}t_6.$$

Manin's conjecture [FMT89] predicts that $N_{U,H}(B)$ grows as $cB(\log B)^{k-1}$ for $B \to \infty$ where $k$ is the rank of the Picard group of the minimal desingularization $\widetilde{S}$ of $S$. As $S$ is a del Pezzo surface of degree 5 whose lines are defined over $\mathbb{Q}$, we have $k = 5$, so our result agrees with this conjecture.

Peyre [Pey95] predicts that $c$ is a product of a constant $\alpha(\widetilde{S})$ whose value is 1/864 by [Der07] and [DJT07] and a product of local densities. We expect that $(1-1/p)^5(1+5/p+1/p^2)$ agrees with the density at each prime $p$, and that $\omega_\infty$ agrees with the real density, but we do not check this here.

Note that $S$ is neither toric nor an equivariant compactification of $\mathbb{G}_\mathrm{a}^2$, so our theorem is not a consequence of [BT98] or [CLT02].

For the proof of the theorem, we use the basic strategy of [BB07], [BBD07] and [DT07] together with the techniques introduced in [BD07]. In Section 2, we translate the counting problem to the question of integral points on a universal torsor and split their counting into three parts. As outlined at the end of Section 2, these parts are handled separately in Sections 3 to 7 and put together again in Section 8 to complete the proof of the theorem.

## 2. A UNIVERSAL TORSOR

We use the notation

$$\boldsymbol{\eta} = (\eta_1, \dots, \eta_4), \quad \boldsymbol{\eta}' = (\eta_1, \dots, \eta_6), \quad \boldsymbol{\alpha} = (\alpha_1, \alpha_2)$$

and, for $(n_1, \dots, n_4) \in \mathbb{Q}^4$,

$$\boldsymbol{\eta}^{(n_1,n_2,n_3,n_4)} := \eta_1^{n_1}\eta_2^{n_2}\eta_3^{n_3}\eta_4^{n_4}.$$

By the method of [DT07] and using the data of [Der06] on the geometry of $S$ and its minimal desingularization $\widetilde{S}$, we obtain a bijection $\Psi : \mathcal{T} \to U(\mathbb{Q})$ with

$$\mathcal{T} := \{(\boldsymbol{\eta}', \boldsymbol{\alpha}) \in \mathbb{Z}^5 \times \mathbb{Z}_{>0} \times \mathbb{Z}^2 \mid \text{(2.1) and coprimality conditions hold}\}$$

where

$$\eta_4\eta_5^2\eta_6 + \eta_1\alpha_1 + \eta_2\alpha_2 = 0 \tag{2.1}$$

and the coprimality conditions are described by the extended Dynkin diagram of $E_1, \dots, E_6, A_1, A_2$ in Figure 1, using the rule that two variables are coprime unless the corresponding divisors in the diagram are connected by an edge. The map $\Psi$ sends $(\boldsymbol{\eta}', \boldsymbol{\alpha}) \in \mathcal{T}$ to

$$(\boldsymbol{\eta}^{(2,2,3,2)}\eta_5, \boldsymbol{\eta}^{(2,1,2,1)}\alpha_1, \eta_6\alpha_1\alpha_2, \boldsymbol{\eta}^{(1,0,1,1)}\eta_5\eta_6\alpha_1, \boldsymbol{\eta}^{(1,2,2,1)}\alpha_2, \boldsymbol{\eta}^{(0,1,1,1)}\eta_5\eta_6\alpha_2)$$

in $U(\mathbb{Q})$.



FIGURE 1. Configuration of curves on $\widetilde{S}$.

Note that these coprimality conditions imply that the formula above for $\Psi(\boldsymbol{\eta}', \alpha)$ results in relatively coprime coordinates $\Psi(\boldsymbol{\eta}', \alpha)_i$, so

$$H(\Psi(\boldsymbol{\eta}', \alpha)) = \max_i\{|\Psi(\boldsymbol{\eta}', \alpha)_i|\}.$$

With (2.1), $H(\Psi(\boldsymbol{\eta}', \boldsymbol{\alpha})) \leqslant B$ implies

$$\boldsymbol{\eta}^{(1,1,2,2)}\eta_5^2|\eta_6| \leqslant 2B, \quad \boldsymbol{\eta}^{(0,0,1,2)}\eta_5^3|\eta_6|^2 \leqslant 2B. \tag{2.2}$$

Using (2.1), the coprimality conditions can be rewritten as

$$\gcd(\alpha_2, \eta_3\eta_5) = 1, \tag{2.3}$$

$$\gcd(\alpha_1, \eta_3\eta_4) = 1, \tag{2.4}$$

$$\gcd(\eta_6, \eta_1\eta_2\eta_3\eta_4) = 1, \tag{2.5}$$

$$\gcd(\eta_5, \eta_1\eta_2\eta_3) = 1, \tag{2.6}$$

$$\gcd(\eta_1, \eta_2) = 1, \ \gcd(\eta_1, \eta_4) = 1, \ \gcd(\eta_2, \eta_4) = 1. \tag{2.7}$$

Therefore, the number $N_{U,H}(B)$ coincides with the number of $(\boldsymbol{\eta}', \boldsymbol{\alpha}) \in \mathbb{Z}_{>0}^5 \times \mathbb{Z}_{\neq 0} \times \mathbb{Z}^2$ which satisfy the torsor equation (2.1), the coprimality conditions (2.3)–(2.7) and the height condition $H(\Psi(\boldsymbol{\eta}', \boldsymbol{\alpha})) \leqslant B$.

Our further strategy is as follows. For fixed $\boldsymbol{\eta}'$, we estimate the number of $\boldsymbol{\alpha}$ satisfying the torsor equation, the coprimality conditions and the height condition. We sum this number over all suitable $\boldsymbol{\eta}'$ afterwards. To get a hold of the error terms in these summations, it will be useful to do this summations in different orders depending on the relative size of $\eta_1, \dots, \eta_6$.

We denote the number of $(\boldsymbol{\eta}', \boldsymbol{\alpha})$ contributing to $N_{U,H}(B)$ that fulfill

$$|\eta_5| \geqslant |\eta_6| \tag{2.8}$$

by $N_a(B)$, and the number of those satisfying

$$|\eta_5| < |\eta_6|. \tag{2.9}$$

by $N_b(B)$.

We split the elements contributing to $N_b(B)$ further into two subsets: For some $A > 0$ to be chosen in Section 8, let $N_{b_1}(B; A)$ be the number of $(\boldsymbol{\eta}', \boldsymbol{\alpha})$ satisfying (2.9) and

$$\boldsymbol{\eta}^{(2,2,3,2)} \leqslant \frac{B}{(\log B)^A}, \tag{2.10}$$

while $N_{b_2}(B; A)$ is the number of the remaining ones.

We deal with $N_{b_2}(B;A)$ in the following Section 3. As a first step for both $N_a(B)$ and $N_{b_1}(B;A)$, we estimate the number of $\boldsymbol{\alpha}$ in Section 5. For $N_a$, we sum first over the bigger $\eta_5$ and then over $\eta_6$ in Section 6, while for $N_{b_1}(B;A)$, we sum in the reverse order in Section 7. The resulting main terms are put together and summed over the remaining variables $\eta_1,\dots,\eta_4$ in Section 8 to complete the proof of the theorem.

## 3. ESTIMATING $N_{b_2}(B;A)$

Our strategy is to estimate the number of $(\boldsymbol{\eta}',\alpha)$ lying in dyadic intervals first, and to sum over all possible intervals in a second step.

**Lemma 1.** *We have* $N_{b_2}(B;A) \ll_A B(\log B)^3(\log\log B)^2$.

*Proof.* Let $\mathcal{N} = \mathcal{N}(N_1,\dots,N_6,A_1,A_2)$ be the number of $(\boldsymbol{\eta}',\boldsymbol{\alpha})$ subject to $N_i/2 < |\eta_i| \leqslant N_i$ for $i \in \{1,\dots,6\}$ and $A_j/2 < |\alpha_j| \leqslant A_j$ for $j \in \{1,2\}$.

Because of the height conditions and using the notation $\mathbf{N}^{(n_1,n_2,n_3,n_4)} := N_1^{n_1}N_2^{n_2}N_3^{n_3}N_4^{n_4}$, we have, if $\mathcal{N} > 0$,

$$B(\log B)^{-A} \ll \mathbf{N}^{(2,2,3,2)} \ll B, \tag{3.1}$$

$$N_6A_1A_2 \ll B, \tag{3.2}$$

$$\mathbf{N}^{(1,0,1,1)}N_5N_6A_1 \ll B, \tag{3.3}$$

$$\mathbf{N}^{(0,1,1,1)}N_5N_6A_2 \ll B, \tag{3.4}$$

$$\mathbf{N}^{(1,1,2,2)}N_5^2N_6 \ll B, \tag{3.5}$$

$$N_5 \ll (\log B)^A. \tag{3.6}$$

Here, (3.5) follows from (2.2). As in [BD07, Lemma 5, 6], we obtain by estimating the number of $\alpha_1,\alpha_2$ in two ways first and summing over $\eta_1,\dots,\eta_6$ afterwards:

$$\mathcal{N} \ll N_3N_4N_5N_6(N_1A_1)^{1/2}(N_2A_2)^{1/2} + N_1N_2N_3N_4N_5N_6.$$

Next, we sum this estimate for $\mathcal{N}(N_1,\dots,N_6,A_1,A_2)$ over all possible dyadic intervals, with $N_1,\dots,N_6,A_1,A_2$ subject to (3.1)–(3.6).

For the first term, we have using (3.2)–(3.5)

$$\begin{aligned}
&\sum_{N_1,\dots,N_6,A_1,A_2} \mathbf{N}^{(1/2,1/2,1,1)}N_5N_6A_1^{1/2}A_2^{1/2}\\
&\ll B^{1/4}\sum_{N_1,\dots,N_6,A_1,A_2} \mathbf{N}^{(1/2,1/2,1,1)}N_5N_6^{3/4}A_1^{1/4}A_2^{1/4}\\
&\ll B^{3/4}\sum_{N_1,\dots,N_6} \mathbf{N}^{(1/4,1/4,1/2,1/2)}N_5^{1/2}N_6^{1/4}\\
&\ll B\sum_{N_1,\dots,N_5} 1\\
&\ll_A B(\log B)^3(\log\log B)^2.
\end{aligned}$$

Here we have used that for fixed $N_2,N_3,N_4$, there are only $O_A(\log\log B)$ possibilities for $N_1$ and $N_5$ by (3.1) and (3.6).

For the second term, we use (3.5) to obtain

$$\begin{aligned}\sum_{N_1,\dots,N_6,A_1,A_2} \mathbf{N}^{(1,1,1,1)} N_5 N_6 &\ll B \sum_{N_1,\dots,N_5,A_1,A_2} \frac{1}{N_3 N_4 N_5} \\ &\ll_A B(\log B)^3 (\log\log B),\end{aligned}$$

which completes the proof. □

## 4. Real-valued functions

Let

$$h(t_0,t_1,t_5,t_6) := \max\left\{\begin{matrix} |t_0^4 t_5|, |t_0^4 t_1|, |t_1 t_5^2 t_6^2 + t_0^2 t_1^2 t_6|, |t_0^2 t_1 t_5 t_6|, \\ |t_0^2 t_5^2 t_6 + t_0^4 t_1|, |t_5^3 t_6^2 + t_0^2 t_1 t_5 t_6| \end{matrix}\right\}. \tag{4.1}$$

Defining

$$Y_0 := \left(\frac{\boldsymbol{\eta}^{(2,2,3,2)}}{B}\right)^{1/5}, \qquad Y_1 := \left(\frac{B}{\boldsymbol{\eta}^{(2,-3,-2,-3)}}\right)^{1/5},$$

$$Y_5 := Y_0^{-1}, \qquad Y_6 := \left(\frac{B}{\boldsymbol{\eta}^{(-3,-3,-2,2)}}\right)^{1/5},$$

we note that the height condition $H(\Psi(\boldsymbol{\eta}', \alpha)) \leqslant B$ is equivalent to

$$h(Y_0, \alpha_1/Y_1, \eta_5/Y_5, \eta_6/Y_6) \leqslant 1.$$

Define

$$g_0(t_0,t_5,t_6) := \int_{h(t_0,t_1,t_5,t_6)\leqslant 1} 1 \,\mathrm{d}t_1, \tag{4.2}$$

$$g_1^a(t_0,t_6;\boldsymbol{\eta};B) := \int_{Y_5 t_5 \geqslant |Y_6 t_6|, t_5>0} g_0(t_0,t_5,t_6)\,\mathrm{d}t_5, \tag{4.3}$$

$$g_1^b(t_0,t_5;\boldsymbol{\eta};B) := \int_{|Y_6 t_6| > \max\{Y_5 t_5, 1\}} g_0(t_0,t_5,t_6)\,\mathrm{d}t_6, \tag{4.4}$$

$$g_2^a(t_0;\boldsymbol{\eta};B) := \int_{|Y_6 t_6|>1} g_1^a(t_0,t_6;\boldsymbol{\eta};B)\,\mathrm{d}t_6, \tag{4.5}$$

$$g_2^b(t_0;\boldsymbol{\eta};B) := \int_0^\infty g_1^b(t_0,t_5;\boldsymbol{\eta};B)\,\mathrm{d}t_5. \tag{4.6}$$

We have

$$\begin{aligned} g_2(t_0;\boldsymbol{\eta};B) &:= g_2^a(t_0;\boldsymbol{\eta};B) + g_2^b(t_0;\boldsymbol{\eta};B) \\ &= \int_{h(t_0,t_1,t_5,t_6)\leqslant 1, |Y_6 t_6|>1, t_5>0} \mathrm{d}t_1\,\mathrm{d}t_5\,\mathrm{d}t_6. \end{aligned} \tag{4.7}$$

**Lemma 2.** *Let $\boldsymbol{\eta} \in \mathbb{Z}_{>0}^4$ be given. Then we have:*

1. $g_0(t_0,t_5,t_6) \ll \frac{1}{t_0|t_6|^{1/2}}$.
2. $g_1^a(t_0,t_6;\boldsymbol{\eta};B) \ll \int_0^\infty g_0(t_0,t_5,t_6)\,\mathrm{d}t_5 \ll \min\{\frac{1}{t_0^{1/2}|t_6|^{5/4}}, \frac{1}{t_0^8}\}$.
3. $g_1^b(t_0,t_5;\boldsymbol{\eta};B) \ll \int_{-\infty}^\infty g_0(t_0,t_5,t_6)\,\mathrm{d}t_6 \ll \frac{1}{t_0 t_5^{3/4}}$.

*Proof.* Since $h(t_0,t_1,t_5,t_6) \leqslant 1$ implies $t_1 \leqslant t_0^{-4}$ and $t_5 \leqslant t_0^{-4}$, the second bound of (2) holds.

It is not hard to check that given $a,b \in \mathbb{R}\backslash\{0\}$, the condition $|at_1^2+bt_1| \leqslant 1$ describes a set of $t_1$ whose length is $\ll |a|^{-1/2}$ for $b^2 \leqslant 8|a|$, while its length is $\ll |b|^{-1} \ll |a|^{-1/2}$ for $b^2 > 8|a|$.

We apply this for $a = t_0^2 t_6$ and $b = t_5^2 t_6^2$, which gives $g_0(t_0,t_5,t_6) \ll (t_0^2|t_6|)^{-1/2}$ which is (1). Integrating it over $t_6 \ll t_5^{3/2}$ (which holds since $|t_0^2 t_1 t_5 t_6| \leqslant 1$ and $|t_5^3 t_6^2 + t_0^2 t_1 t_5 t_6| \leqslant 1$ imply $|t_5^3 t_6^2| \leqslant 2$) results in (3).

For the first bound of (2), we distinguish the case $t_5^4 t_6^4 \leqslant 8t_0^2|t_6|$ and its opposite. In the first case, we combine $t_5 \ll t_0^{1/2}|t_6|^{-3/4}$ with (1). In the second case, we integrate $g_0(t_0,t_5,t_6) \ll t_5^{-2}|t_6|^{-2}$ over $t_5 \gg t_0^{1/2}|t_6|^{-3/4}$. □

Finally, we define

$$G_2(t_0) := \int_{h(t_0,t_1,t_5,t_6)\leqslant 1, t_5>0} \mathrm{d}t_1 \, \mathrm{d}t_5 \, \mathrm{d}t_6 \tag{4.8}$$

which is related to $\omega_\infty$ defined in the statement of our theorem:

**Lemma 3.** *For any $t_0 > 0$, we have $G_2(t_0) = \frac{\omega_\infty}{t_0^2}$.*

*Proof.* Similar to [BD07, Lemma 7]. □

## 5. ESTIMATING $N_a(B)$ AND $N_{b_1}(B;A)$ – FIRST STEP

For fixed $\boldsymbol{\eta}'$ subject to the coprimality conditions (2.5)–(2.7), let $N_0$ be the number of $\alpha_1, \alpha_2$ subject to (2.1), $h(Y_0, \alpha_1/Y_1, \eta_5/Y_5, \eta_6/Y_6) \leqslant 1$ and the coprimality conditions (2.3), (2.4).

We remove (2.3) by a Möbius inversion and obtain

$$N_0 = \sum_{k_2|\eta_3\eta_5} \mu(k_2) \# \left\{ \alpha_1 \;\middle|\; \begin{array}{l} \eta_4\eta_5^2\eta_6 \equiv -\eta_1\alpha_1 \pmod{k_2\eta_2}, \\ h(Y_0,\alpha_1/Y_1,\eta_5/Y_5,\eta_6/Y_6) \leqslant 1, \\ \text{(2.4) holds} \end{array} \right\}.$$

The summand vanishes unless $\gcd(k_2, \eta_1\eta_4) = 1$. Since $\eta_3, \eta_5$ are coprime, we write $k_2 = k_{23}k_{25}$ uniquely such that $k_{2i} \mid \eta_i$ for $i \in \{3,5\}$. We check that $k_{25}|\alpha_1$. We write $\eta_5 = k_{25}\eta_5'$, $\alpha_1 = k_{25}\alpha_1'$ and obtain

$$N_0 = \sum_{\substack{k_{23}|\eta_3, k_{25}|\eta_5 \\ \gcd(k_{23}k_{25},\eta_1\eta_4)=1}} \mu(k_{23})\mu(k_{25}) N_0(k_{23},k_{25})$$

where

$$N_0(k_{23},k_{25}) = \# \left\{ \alpha_1' \;\middle|\; \begin{array}{l} k_{25}\eta_4\eta_5'^2\eta_6 \equiv -\eta_1\alpha_1' \pmod{k_{23}\eta_2}, \\ h(Y_0,\alpha_1' k_{25}/Y_1,\eta_5/Y_5,\eta_6/Y_6) \leqslant 1, \\ \gcd(k_{25}\alpha_1',\eta_3\eta_4) = 1 \end{array} \right\}.$$

Note that $\gcd(k_{25},\eta_3\eta_4) = 1$ holds automatically, so we may remove this condition. We remove the coprimality condition for $\alpha_1'$ by another Möbius inversion and obtain, writing $\alpha_1' = k_1\alpha_1''$,

$$N_0(k_{23},k_{25}) = \sum_{k_1|\eta_3\eta_4} \mu(k_1) \# \left\{ \alpha_1'' \;\middle|\; \begin{array}{l} k_{25}\eta_4\eta_5'^2\eta_6 \equiv -k_1\eta_1\alpha_1'' \pmod{k_{23}\eta_2}, \\ h(Y_0,\alpha_1'' k_{25}k_1/Y_1,\eta_5/Y_5,\eta_6/Y_6) \leqslant 1, \end{array} \right\}.$$

Note that the summand vanishes unless $\gcd(k_1, k_{23}\eta_2) = 1$, so we may restrict the summation over $k_1 | \eta_3\eta_4$ subject to $\gcd(k_1, k_{23}\eta_2) = 1$. Since then $\gcd(k_1\eta_1, k_{23}\eta_2) = 1$, the number of $\alpha_1''$ is

$$\frac{Y_1}{k_1 k_{23} k_{25} \eta_2} g_0(Y_0, \eta_5/Y_5, \eta_6/Y_6) + O(1).$$

Define $\phi^*(n) := \prod_{p|n}(1 - 1/p)$.

**Lemma 4.** *We have*

$$N_0 = \frac{Y_1}{\eta_2} g_0(Y_0, \eta_5/Y_5, \eta_6/Y_6)\vartheta_0(\boldsymbol{\eta})\frac{\phi^*(\eta_5)}{\phi^*(\gcd(\eta_5, \eta_4))} + O(R_0(\boldsymbol{\eta}, \eta_5, \eta_6))$$

*with*

$$\vartheta_0(\boldsymbol{\eta}) := \sum_{\substack{k_{23}|\eta_3 \\ \gcd(k_{23}, \eta_1\eta_4)=1}} \frac{\mu(k_{23})\phi^*(\eta_3\eta_4)}{k_{23}\phi^*(\gcd(\eta_3, k_{23}\eta_2))}$$

*and*

$$\sum_{\eta_1,\dots,\eta_6} R_0(\boldsymbol{\eta}, \eta_5, \eta_6) \ll B(\log B)^2.$$

*Proof.* For the main term, note that

$$\begin{aligned} &\sum_{\substack{k_{23}|\eta_3, k_{25}|\eta_5 \\ \gcd(k_{23}k_{25}, \eta_1\eta_4)=1}} \frac{\mu(k_{23})\mu(k_{25})}{k_{23}k_{25}} \sum_{\substack{k_1|\eta_3\eta_4 \\ \gcd(k_1, k_{23}\eta_2)=1}} \frac{\mu(k_1)}{k_1} \\ = &\sum_{\substack{k_{23}|\eta_3 \\ \gcd(k_{23}, \eta_1\eta_4)=1}} \frac{\mu(k_{23})}{k_{23}} \cdot \frac{\phi^*(\eta_5)}{\phi^*(\gcd(\eta_5, \eta_1\eta_4))} \cdot \frac{\phi^*(\eta_3\eta_4)}{\phi^*(\gcd(\eta_3\eta_4, k_{23}\eta_2))}. \end{aligned}$$

Using $\gcd(\eta_5, \eta_1) = 1$ and $\gcd(\eta_4, k_{23}\eta_2) = 1$, we obtain $\vartheta_0$.

We have

$$R_0(\boldsymbol{\eta}, \eta_5, \eta_6) \ll 2^{\omega(\eta_3)+\omega(\eta_5)+\omega(\eta_3\eta_4)}.$$

We sum this over all suitable $\eta_1, \dots, \eta_6$ and use (2.2) to obtain

$$\begin{aligned} \sum_{\eta_1,\dots,\eta_6} R_0(\boldsymbol{\eta}, \eta_5, \eta_6) &\ll \sum_{\eta_1,\dots,\eta_5} \frac{2^{\omega(\eta_3)+\omega(\eta_5)+\omega(\eta_3\eta_4)} B}{\boldsymbol{\eta}^{(1,1,2,2)}\eta_5^2} \\ &\ll B(\log B)^2, \end{aligned}$$

completing the proof of this lemma. □

## 6. ESTIMATING $N_a(B)$ – SECOND STEP

Let $N_1^a := N_1^a(\boldsymbol{\eta}, \eta_6; B)$ be the sum of the main term of Lemma 4 over $\eta_5$ subject to (2.6) and (2.8). We sum the main term of $N_1^a$ over $\eta_6$ afterwards to obtain $N_2^a(\boldsymbol{\eta}; B)$.

Using [BD07, Lemma 2] with $\alpha = 0$, $q = 1$, we obtain (where $f_{a,b}(n)$ is defined to be $\phi^*(n)/\phi^*(\gcd(n,a))$ if $\gcd(n,b) = 1$ and to be zero otherwise)

$$\begin{aligned} N_1^a &= \frac{Y_1}{\eta_2}\vartheta_0(\boldsymbol{\eta}) \sum_{\eta_5 \geqslant |\eta_6|} f_{\eta_4,\eta_1\eta_2\eta_3}(\eta_5) g_0(Y_0, \eta_5/Y_5, \eta_6/Y_6; \boldsymbol{\eta}; B) \\ &= \frac{Y_1 Y_5}{\eta_2} g_1^a(Y_0, \eta_6/Y_6; \boldsymbol{\eta}; B)\vartheta_0(\boldsymbol{\eta}) \frac{\phi^*(\eta_1\eta_2\eta_3)}{\zeta(2)} \prod_{p \mid \eta_1\eta_2\eta_3\eta_4} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &\quad + O\left(\frac{Y_1}{\eta_2}|\vartheta_0(\boldsymbol{\eta})|(\log B) 2^{\omega(\eta_1\eta_2\eta_3)} \sup_{t_5} g_0(Y_0, t_5, \eta_6/Y_6; \boldsymbol{\eta}; B)\right), \end{aligned}$$

where the supremum is taken over $t_5 \geqslant |\eta_6|/Y_5$.

**Lemma 5.** *We have*

$$N_1^a = \frac{Y_1 Y_5}{\eta_2} g_1^a(Y_0, \eta_6/Y_6; \boldsymbol{\eta}; B)\vartheta_1^a(\boldsymbol{\eta}) + O(R_1^a(\boldsymbol{\eta}, \eta_6; B))$$

*with*

$$\vartheta_1^a(\boldsymbol{\eta}) := \vartheta_0(\boldsymbol{\eta}) \frac{\phi^*(\eta_1\eta_2\eta_3)}{\zeta(2)} \prod_{p \mid \eta_1\eta_2\eta_3\eta_4} \left(1 - \frac{1}{p^2}\right)^{-1}$$

*and*

$$\sum_{\boldsymbol{\eta}, \eta_6} R_1^a(\boldsymbol{\eta}, \eta_6; B) \ll B \log B.$$

*Proof.* The main term is clear. Define $\phi^\dagger(n) := \prod p|n(1 + 1/p)$. For the error term, we use Lemma 2(1) to estimate its sum over $\boldsymbol{\eta}, \eta_6$ as

$$\begin{aligned} &\ll \sum_{\boldsymbol{\eta}, \eta_6} \frac{2^{\omega(\eta_1\eta_2\eta_3)}\phi^\dagger(\eta_3) Y_1 Y_6^{1/2} \log B}{Y_0 \eta_2 |\eta_6|^{1/2}} \\ &= \sum_{\boldsymbol{\eta}, \eta_6} \frac{2^{\omega(\eta_1\eta_2\eta_3)}\phi^\dagger(\eta_3) B^{1/2} \log B}{\eta_1^{1/2}\eta_2^{1/2}|\eta_6|^{1/2}} \\ &\ll \sum_{\eta_1,\eta_2,\eta_3,\eta_6} \frac{2^{\omega(\eta_1\eta_2\eta_3)}\phi^\dagger(\eta_3) B \log B}{\boldsymbol{\eta}^{(5/4,5/4,5/4,0)}|\eta_6|^{3/2}} \\ &\ll B \log B. \end{aligned}$$

Here, we use

$$\eta_4 \leqslant \left(\frac{B}{\boldsymbol{\eta}^{(2,2,3,0)}|\eta_6|}\right)^{1/4} \cdot \left(\frac{B}{\boldsymbol{\eta}^{(1,1,2,0)}|\eta_6|^3}\right)^{1/4} = \frac{B^{1/2}}{\boldsymbol{\eta}^{(3/4,3/4,5/4,0)}|\eta_6|}$$

which is obtained with (2.8). $\square$

To sum the main term of $N_1^a$ over $\eta_6$, we remove the coprimality condition (2.5) by a Möbius inversion and obtain, writing $\eta_6 = k_6\eta_6'$ and applying partial summation,

$$\begin{aligned} N_2^a &= \frac{Y_1Y_5}{\eta_2}\vartheta_1^a(\boldsymbol{\eta}) \sum_{k_6 \mid \eta_1\eta_2\eta_3\eta_4} \mu(k_6) \sum_{|\eta_6'| \geqslant 1} g_1^a(Y_0, \eta_6' k_6/Y_6; \boldsymbol{\eta}; B) \\ &= \frac{Y_1Y_5Y_6}{\eta_2}\vartheta_1^a(\boldsymbol{\eta}) \sum_{k_6 \mid \eta_1\eta_2\eta_3\eta_4} \frac{\mu(k_6)}{k_6} \int_{|t_6| \geqslant k_6/Y_6} g_1^a(Y_0, t_6; \boldsymbol{\eta}; B)\,\mathrm{d}t_6 \\ &\quad + O\left( \frac{Y_1Y_5}{\eta_2} |\vartheta_1^a(\boldsymbol{\eta})| \sum_{k_6 \mid \eta_1\eta_2\eta_3\eta_4} |\mu(k_6)| \sup_{|t_6| \geqslant k_6/Y_6} g_1^a(Y_0, t_6; \boldsymbol{\eta}; B) \right). \end{aligned}$$

**Lemma 6.** *We have*

$$N_2^a = \frac{Y_1Y_5Y_6}{\eta_2} g_2^a(Y_0; \boldsymbol{\eta}; B)\vartheta_2^a(\boldsymbol{\eta}) + O(R_2^a(\boldsymbol{\eta}; B))$$

*with*

$$\vartheta_2^a(\boldsymbol{\eta}) := \vartheta_1^a(\boldsymbol{\eta})\phi^*(\eta_1\eta_2\eta_3\eta_4)$$

*and*

$$\sum_{\boldsymbol{\eta}} R_2^a(\boldsymbol{\eta}; B) \ll B(\log B)^{4-1/5}.$$

*Proof.* In order to replace the integral over $|t_6| \geqslant k_6/Y_6$ in the estimation before the statement of the lemma by $g_2^a(Y_0; \boldsymbol{\eta}; B)$, we must add

$$\frac{Y_1Y_5Y_6}{\eta_2}\vartheta_1^a(\boldsymbol{\eta}) \sum_{k_6 \mid \eta_1\eta_2\eta_3\eta_4} \frac{\mu(k_6)}{k_6} \int_{1/Y_6 < |t_6| < k_6/Y_6} g_1^a(Y_0, t_6; \boldsymbol{\eta}; B)\,\mathrm{d}t_6$$

as a second error term.

We distinguish the case

$$\boldsymbol{\eta}^{(3,3,4,2)} < \lambda B \tag{6.1}$$

for some $\lambda > 0$ to be chosen later, giving a total contribution $E_1(\lambda)$ to the error term, and its opposite

$$\boldsymbol{\eta}^{(3,3,4,2)} \geqslant \lambda B, \tag{6.2}$$

contributing in total $E_2(\lambda)$.

Starting with $E_1(\lambda)$, we use the first bound of Lemma 2(2). For the first error term, we obtain

$$\begin{aligned} &\ll \sum_{\boldsymbol{\eta}} \sum_{k_6 \mid \eta_1\eta_2\eta_3\eta_4} \frac{|\mu(k_6)|\phi^\dagger(\eta_3)Y_1Y_5Y_6^{5/4}}{k_6^{5/4}\eta_2 Y_0^{1/2}} \\ &\ll \sum_{\boldsymbol{\eta}} \frac{\phi^\dagger(\eta_3)B^{3/4}}{\boldsymbol{\eta}^{(1/4,1/4,0,1/2)}} \\ &\ll \sum_{\eta_1,\eta_2,\eta_3} \frac{\phi^\dagger(\eta_3)\lambda^{1/4}B}{\boldsymbol{\eta}^{(1,1,1,0)}} \\ &\ll \lambda^{1/4}B(\log B)^3. \end{aligned}$$

For the second error term, we use

$$\int_{1/Y_6}^{k_6/Y_6} g_1^a(Y_0,t_6;\boldsymbol{\eta};B)\,\mathrm{d}t_6 \ll \int_{1/Y_6}^{k_6/Y_6} \frac{1}{Y_0^{1/2}|t_6|^{5/4}}\,\mathrm{d}t_6 \ll \frac{Y_6^{1/4}}{Y_0^{1/2}}$$

and obtain

$$\begin{aligned}
&\ll \sum_{\boldsymbol{\eta}} \sum_{k_6|\eta_1\eta_2\eta_3\eta_4} \frac{|\mu(k_6)|\phi^\dagger(\eta_3)Y_1Y_5Y_6^{5/4}}{k_6\eta_2Y_0^{1/2}}\\
&\ll \sum_{\boldsymbol{\eta}} \frac{\phi^\dagger(\eta_1\eta_2\eta_3\eta_4)\phi^\dagger(\eta_3)B^{3/4}}{\boldsymbol{\eta}^{(1/4,1/4,0,1/2)}}\\
&\ll \sum_{\eta_1,\eta_2,\eta_3} \frac{\phi^\dagger(\eta_1\eta_2\eta_3)\phi^\dagger(\eta_3)\lambda^{1/4}B}{\boldsymbol{\eta}^{(1,1,1,0)}}\\
&\ll \lambda^{1/4}B(\log B)^3.
\end{aligned}$$

Therefore, $E_1(\lambda) \ll \lambda^{1/4}B(\log B)^3$.

For $E_2(\lambda)$, we use the second bound of Lemma 2(3). For the first part of this error term, we get

$$\begin{aligned}
&\ll \sum_{\boldsymbol{\eta}} \sum_{k_6|\eta_1\eta_2\eta_3\eta_4} \frac{|\mu(k_6)|\phi^\dagger(\eta_3)Y_1Y_5}{\eta_2Y_0^8}\\
&\ll \sum_{\boldsymbol{\eta}} \frac{2^{\omega(\eta_1\eta_2\eta_3\eta_4)}\phi^\dagger(\eta_3)B^2}{\boldsymbol{\eta}^{(4,4,5,3)}}\\
&\ll \sum_{\eta_1,\eta_2,\eta_3} \frac{2^{\omega(\eta_1\eta_2\eta_3)}\phi^\dagger(\eta_3)B\log B}{\lambda\boldsymbol{\eta}^{(1,1,1,0)}}\\
&\ll \lambda^{-1}B(\log B)^7.
\end{aligned}$$

For the second part of the error term, we use

$$\int_{1/Y_6}^{k_6/Y_6} g_1^a(Y_0,t_6;\boldsymbol{\eta};B)\,\mathrm{d}t_6 \ll \int_{1/Y_6}^{k_6/Y_6} \frac{1}{Y_0^8}\,\mathrm{d}t_6 \ll \frac{k_6}{Y_0^8Y_6}$$

and obtain

$$\begin{aligned}
&\ll \sum_{\boldsymbol{\eta}} \sum_{k_6|\eta_1\eta_2\eta_3\eta_4} \frac{|\mu(k_6)|\phi^\dagger(\eta_3)Y_1Y_5Y_6}{k_6\eta_2}\cdot\frac{k_6}{Y_0^8Y_6}\\
&\ll \sum_{\boldsymbol{\eta}} \frac{2^{\omega(\eta_1\eta_2\eta_3\eta_4)}\phi^\dagger(\eta_3)B^2}{\boldsymbol{\eta}^{(4,4,5,3)}}\\
&\ll \sum_{\eta_1,\eta_2,\eta_3} \frac{2^{\omega(\eta_1\eta_2\eta_3)}\phi^\dagger(\eta_3)B\log B}{\lambda\boldsymbol{\eta}^{(1,1,1,0)}}\\
&\ll \lambda^{-1}B(\log B)^7.
\end{aligned}$$

In total, $E_2(\lambda) \ll \lambda^{-1}B(\log B)^7$.

Choosing $\lambda = (\log B)^{16/5}$ gives a total error term of $O(B(\log B)^{4-1/5})$. $\square$

## 7. ESTIMATING $N_{b_1}(B;A)$ – SECOND STEP

Let $N_1^b := N_1^b(\boldsymbol{\eta}, \eta_5; B)$ be the main term of $N_0$ in Lemma 4 summed over $\eta_6$ subject to (2.5) and (2.9). We denote the main term of this summed over all $\eta_5$ by $N_2^b := N_2^b(\boldsymbol{\eta}; B)$.

We remove (2.5) by a Möbius inversion and get

$$N_1^b = \frac{Y_1}{\eta_2}\vartheta_0(\boldsymbol{\eta})\frac{\phi^*(\eta_5)}{\phi^*(\gcd(\eta_5,\eta_4))}\sum_{k_6 | \eta_1\eta_2\eta_3\eta_4} \mu(k_6)A$$

where

$$A = \sum_{\substack{\eta_6' \in \mathbb{Z}_{\neq 0} \\ k_6|\eta_6'| > \eta_5}} g_0(Y_0, \eta_5/Y_5, k_6\eta_6'/Y_6; \boldsymbol{\eta}; B).$$

By partial summation,

$$A = \frac{Y_6}{k_6} g_1^b(Y_0, \eta_5/Y_5; \boldsymbol{\eta}; B) + O(\sup_{t_6} g_0(Y_0, \eta_5/Y_5, t_6)),$$

where the supremum is taken over $t_6$ subject to $|t_6| > \eta_5/Y_6$.

**Lemma 7.** *We have*

$$N_1^b = \frac{Y_1 Y_6}{\eta_2} g_1^b(Y_0, \eta_5/Y_5; \boldsymbol{\eta}; B)\vartheta_1^b(\boldsymbol{\eta})\frac{\phi^*(\eta_5)}{\phi^*(\gcd(\eta_5,\eta_4))} + O(R_1^b(\boldsymbol{\eta}, \eta_5; B))$$

*with*

$$\vartheta_1^b(\boldsymbol{\eta}) := \vartheta_0(\boldsymbol{\eta})\phi^*(\eta_1\eta_2\eta_3\eta_4)$$

*and*

$$\sum_{\boldsymbol{\eta},\eta_5} R_1^b(\boldsymbol{\eta}, \eta_5; B) \ll B \log B.$$

*Proof.* The main term is clear. We apply Lemma 2(1) to deduce that the error term can be estimated as

$$\begin{aligned}
\sum_{\boldsymbol{\eta},\eta_5} R_1^b(\boldsymbol{\eta}, \eta_5; B) &\ll \sum_{\boldsymbol{\eta},\eta_5} \frac{2^{\omega(\eta_1\eta_2\eta_3\eta_4)}\phi^\dagger(\eta_3)Y_1 Y_6^{1/2}}{\eta_2\eta_5^{1/2}Y_0} \\
&= \sum_{\boldsymbol{\eta},\eta_5} \frac{2^{\omega(\eta_1\eta_2\eta_3\eta_4)}\phi^\dagger(\eta_3)B^{1/2}}{\eta_1^{1/2}\eta_2^{1/2}\eta_5^{1/2}} \\
&\ll \sum_{\eta_1,\eta_2,\eta_3,\eta_5} \frac{2^{\omega(\eta_1\eta_2\eta_3)}\phi^\dagger(\eta_3)B\log B}{\boldsymbol{\eta}^{(5/4,5/4,5/4,0)}\eta_5^{3/2}} \\
&\ll B\log B.
\end{aligned}$$

In the last step, we have used

$$\eta_4 \leqslant \left(\frac{B}{\boldsymbol{\eta}^{(2,2,3,0)}\eta_5}\right)^{1/4} \cdot \left(\frac{B}{\boldsymbol{\eta}^{(1,1,2,0)}\eta_5^3}\right)^{1/4} = \frac{B^{1/2}}{\boldsymbol{\eta}^{(3/4,3/4,5/4,0)}\eta_5},$$

which we obtain using (2.2) and (2.9). ☐

Next, we sum the main term of Lemma 5 over all suitable $\eta_5$. Apply [BD07, Lemma 2] with $\alpha = 0$, $q = 1$ to obtain

$$\begin{aligned} N_2^b &= \frac{Y_1Y_6}{\eta_2}\vartheta_1^b(\boldsymbol{\eta}) \sum_{\eta_5 \geqslant 1} f_{\eta_4,\eta_1\eta_2\eta_3}(\eta_5) g_1^b(Y_0, \eta_5/Y_5; \boldsymbol{\eta}; B) \\ &= \frac{Y_1Y_5Y_6}{\eta_2} g_2^b(Y_0; \boldsymbol{\eta}; B)\vartheta_1^b(\boldsymbol{\eta}) \frac{\phi^*(\eta_1\eta_2\eta_3)}{\zeta(2)} \prod_{p \mid \eta_1\eta_2\eta_3\eta_4} \left(1 - \frac{1}{p^2}\right)^{-1} \\ &\quad + O\left(\frac{Y_1Y_6}{\eta_2} |\vartheta_1^b(\boldsymbol{\eta})| (\log B) 2^{\omega(\eta_1\eta_2\eta_3)} \sup_{t_5} g_1^b(Y_0, t_5; \boldsymbol{\eta}; B)\right) \\ &\quad + O\left(\frac{Y_1Y_5Y_6}{\eta_2} |\vartheta_1^b(\boldsymbol{\eta})| \int_{0 \leqslant t_5 \leqslant 1/Y_5} g_1^b(Y_0, t_5; \boldsymbol{\eta}; B)\, \mathrm{d}t_5\right), \end{aligned}$$

where the supremum is taken over $t_5 > 1/Y_5$.

**Lemma 8.** *We have*

$$N_2^b = \frac{Y_1Y_5Y_6}{\eta_2} g_2^b(Y_0; \boldsymbol{\eta}; B)\vartheta_2^b(\boldsymbol{\eta}) + O(R_2^b(\boldsymbol{\eta}; B))$$

*with*

$$\vartheta_2^b(\boldsymbol{\eta}) := \vartheta_1^b(\boldsymbol{\eta}) \frac{\phi^*(\eta_1\eta_2\eta_3)}{\zeta(2)} \prod_{p \mid \eta_1\eta_2\eta_3\eta_4} \left(1 - \frac{1}{p^2}\right)^{-1}$$

*and*

$$\sum_{\boldsymbol{\eta}} R_2^b(\boldsymbol{\eta}; B) \ll B(\log B)^{7-A/4},$$

*where the sum is taken over* $\boldsymbol{\eta}$ *satisfying* (2.10).

*Proof.* The main term is clear from the discussion before the lemma. The first part of the error term makes the contribution

$$\ll \sum_{\boldsymbol{\eta}} \frac{2^{\omega(\eta_1\eta_2\eta_3)} \phi^\dagger(\eta_3) Y_1 Y_6 \log B}{\eta_2} \sup g_1^b(Y_0, t_5; \boldsymbol{\eta}; B).$$

We use Lemma 2(3) and (2.10) to obtain

$$\begin{aligned} &\ll \sum_{\boldsymbol{\eta}} \frac{2^{\omega(\eta_1\eta_2\eta_3)} \phi^\dagger(\eta_3) Y_1 Y_5^{3/4} Y_6 \log B}{\eta_2 Y_0} \\ &= \sum_{\boldsymbol{\eta}} \frac{2^{\omega(\eta_1\eta_2\eta_3)} \phi^\dagger(\eta_3) B^{3/4} \log B}{\boldsymbol{\eta}^{(1/2,1/2,1/4,1/2)}} \\ &\ll \sum_{\eta_1,\eta_2,\eta_3} \frac{2^{\omega(\eta_1\eta_2\eta_3)} \phi^\dagger(\eta_3) B (\log B)^{1-A/4}}{\boldsymbol{\eta}^{(1,1,1,0)}} \\ &\ll B(\log B)^{7-A/4}. \end{aligned}$$

The contribution of the second term is, using Lemma 2(3) and (2.10) again,

$$
\begin{aligned}
&\ll \sum_{\boldsymbol{\eta}} \frac{\phi^\dagger(\eta_3) Y_1 Y_5 Y_6}{\eta_2} \int_0^{1/Y_5} \frac{1}{Y_0 t_5^{3/4}} \,\mathrm{d}t_5 \\
&\ll \sum_{\boldsymbol{\eta}} \frac{\phi^\dagger(\eta_3) Y_1 Y_6^{5/4} Y_6}{\eta_2 Y_0} \\
&= \sum_{\boldsymbol{\eta}} \frac{\phi^\dagger(\eta_3) B^{3/4}}{\boldsymbol{\eta}^{(1/2,1/2,1/4,1/2)}} \\
&\ll \sum_{\eta_1,\eta_2,\eta_3} \frac{\phi^\dagger(\eta_3) B (\log B)^{-A/4}}{\boldsymbol{\eta}^{(1,1,1,0)}} \\
&\ll B(\log B)^{3-A/4}.
\end{aligned}
$$

This completes the proof of the lemma. □

## 8. The final step

By the discussion at the end of Section 2, we have, for any $A > 0$,

$$N_{U,H}(B) = N_a(B) + N_{b_1}(B; A) + N_{b_2}(B; A).$$

By Lemma 1,

$$N_{U,H}(B) = N_a(B) + N_{b_1}(B; A) + O_A(B(\log B)^3 (\log\log B)^2).$$

Using Lemmas 4, 5 and 6 and combining their error terms shows that

$$N_a(B) = \sum_{\boldsymbol{\eta} \in \mathcal{E}(B)} \frac{Y_1 Y_5 Y_6}{\eta_2} g_2^a(Y_0; \boldsymbol{\eta}; B) \vartheta_2^a(\boldsymbol{\eta}) + O(B(\log B)^{4-1/5}),$$

where

$$\mathcal{E}(t) := \{\boldsymbol{\eta} \in \mathbb{Z}_{>0}^4 \mid (2.7), \boldsymbol{\eta}^{(2,2,3,2)} \leqslant t\}$$

for any $t \geqslant 1$, while Lemmas 4, 7 and 8 give, choosing $A := 28$,

$$N_{b_1}(B; 28) = \sum_{\boldsymbol{\eta} \in \mathcal{E}(B/(\log B)^{28})} \frac{Y_1 Y_5 Y_6}{\eta_2} g_2^b(Y_0; \boldsymbol{\eta}; B) \vartheta_2^b(\boldsymbol{\eta}) + O(B(\log B)^2).$$

Recall the definition (4.7) of $g_2$.

**Lemma 9.** *We have*

$$N_{U,H}(B) = \sum_{\boldsymbol{\eta} \in \mathcal{E}(B)} \frac{Y_1 Y_5 Y_6}{\eta_2} g_2(Y_0; \boldsymbol{\eta}; B) \vartheta(\boldsymbol{\eta}) + O(B(\log B)^{4-1/5})$$

*where*

$$
\begin{aligned}
\vartheta(\boldsymbol{\eta}) := {}& \frac{\phi^*(\eta_3\eta_4)\phi^*(\eta_1\eta_2\eta_3)\phi^*(\eta_1\eta_2\eta_3\eta_4)}{\zeta(2)} \prod_{p \mid \eta_1\eta_2\eta_3\eta_4} \left(1 - \frac{1}{p^2}\right)^{-1} \\
& \times \left( \sum_{\substack{k_{23} \mid \eta_3 \\ \gcd(k_{23}, \eta_1\eta_4) = 1}} \frac{\mu(k_{23})}{k_{23}\phi^*(\gcd(\eta_3, k_{23}\eta_2))} \right)
\end{aligned}
$$

*if the coprimality conditions* (2.7) *hold, and* $\vartheta(\boldsymbol{\eta}) := 0$ *otherwise.*

*Proof.* We easily check that $\vartheta(\boldsymbol{\eta})$ agrees with $\vartheta_2^a(\boldsymbol{\eta})$ and $\vartheta_2^b(\boldsymbol{\eta})$ for $\boldsymbol{\eta}$ satisfying (2.7).

In view of the discussion before the lemma, it remains to show that

$$\sum_{\boldsymbol{\eta} \in \mathcal{E}(B) \setminus \mathcal{E}(B/(\log B)^{28})} \frac{Y_1 Y_5 Y_6}{\eta_2} g_2^b(Y_0; \boldsymbol{\eta}; B) \vartheta_2^b(\boldsymbol{\eta})$$

makes a negligible contribution.

Indeed, we estimate this as

$$\begin{aligned} &\ll \sum_{\boldsymbol{\eta} \in \mathcal{E}(B) \setminus \mathcal{E}(B/(\log B)^{28})} \frac{\phi^\dagger(\eta_3) Y_1 Y_5 Y_6}{\eta_2 Y_0^2} \\ &= \sum_{\boldsymbol{\eta} \in \mathcal{E}(B) \setminus \mathcal{E}(B/(\log B)^{28})} \frac{\phi^\dagger(\eta_3) B}{\boldsymbol{\eta}^{(1,1,1,1)}} \\ &\ll B(\log B)^3 (\log \log B) \end{aligned}$$

since we have

$$g_2^b(t_0; \boldsymbol{\eta}; B) \ll \int_0^{1/t_0^4} g_1^b(t_0, t_5; \boldsymbol{\eta}; B) \, \mathrm{d}t_5 \ll \int_0^{1/t_0^4} \frac{1}{t_0 t_5^{3/4}} \, \mathrm{d}t_5 \ll \frac{1}{t_0^2},$$

using Lemma 2(3) and the fact that $g_1^b(t_0, t_5; \boldsymbol{\eta}; B) = 0$ unless $t_5 \ll 1/t_0^4$ by (4.1). $\square$

Define

$$\mathcal{E}^*(B) := \{\boldsymbol{\eta} \in \mathbb{Z}_{>0}^4 \mid \boldsymbol{\eta}^{(2,2,3,2)} \leqslant B, \boldsymbol{\eta}^{(3,3,4,2)} > B\}.$$

**Lemma 10.** *We have*

$$N_{U,H}(B) = \omega_\infty B \sum_{\boldsymbol{\eta} \in \mathcal{E}^*(B)} \frac{\vartheta(\boldsymbol{\eta})}{\boldsymbol{\eta}^{(1,1,1,1)}} + O(B(\log B)^{4-1/5}).$$

*Proof.* By Lemma 2(2), we have

$$g_2(Y_0; \boldsymbol{\eta}; B) \ll \int_{|Y_6 t_6| > 1} \frac{1}{Y_0^{1/2} |t_6|^{5/4}} \, \mathrm{d}t_6 \ll \frac{Y_6^{1/4}}{Y_0^{1/2}}.$$

Therefore,

$$\begin{aligned} \sum_{\boldsymbol{\eta} \in \mathcal{E}(B) \setminus \mathcal{E}^*(B)} \frac{Y_1 Y_5 Y_6}{\eta_2} g_2(Y_0; \boldsymbol{\eta}; B) \vartheta(\boldsymbol{\eta}) &\ll \sum_{\boldsymbol{\eta}^{(3,3,4,2)} \leqslant B} \frac{\phi^\dagger(\eta_3) Y_1 Y_5 Y_6^{5/4}}{\eta_2 Y_0^{1/2}} \\ &\ll \sum_{\boldsymbol{\eta}^{(3,3,4,2)} \leqslant B} \frac{\phi^\dagger(\eta_3) B^{3/4}}{\boldsymbol{\eta}^{(1/4,1/4,0,1/2)}} \\ &\ll \sum_{\eta_1, \eta_2, \eta_3} \frac{\phi^\dagger(\eta_3) B}{\boldsymbol{\eta}^{(1,1,1,0)}} \\ &\ll B(\log B)^3. \end{aligned}$$

This proves that

$$N_{U,H}(B) = \sum_{\boldsymbol{\eta} \in \mathcal{E}^*(B)} \frac{Y_1Y_5Y_6}{\eta_2} g_2(Y_0; \boldsymbol{\eta}; B)\vartheta(\boldsymbol{\eta}) + O(B(\log B)^{4-1/5}).$$

Comparing the definitions (4.7) and (4.8) of the functions $g_2$ and $G$ together with the estimation

$$\begin{aligned} &\sum_{\boldsymbol{\eta} \in \mathcal{E}^*(B)} \frac{Y_1Y_5Y_6}{\eta_2} \int_{h(Y_0,t_1,t_5,t_6) \leqslant 1, |Y_6 t_6| \leqslant 1, t_5 > 0} \mathrm{d}t_1 \, \mathrm{d}t_5 \, \mathrm{d}t_6 \\ \ll &\sum_{\boldsymbol{\eta} \in \mathcal{E}^*(B)} \frac{\phi^\dagger(\eta_3) Y_1Y_5Y_6}{\eta_2} \int_{|Y_6t_6| \leqslant 1} \frac{1}{Y_0^8} \, \mathrm{d}t_6 \\ \ll &\sum_{\boldsymbol{\eta} \in \mathcal{E}^*(B)} \frac{\phi^\dagger(\eta_3) Y_1Y_5}{\eta_2 Y_0^8} \\ = &\sum_{\boldsymbol{\eta} \in \mathcal{E}^*(B)} \frac{\phi^\dagger(\eta_3) B^2}{\boldsymbol{\eta}^{(4,4,5,3)}} \\ \ll &\sum_{\eta_1,\eta_2,\eta_3} \frac{\phi^\dagger(\eta_3) B}{\boldsymbol{\eta}^{(1,1,1,0)}} \\ \ll &\, B(\log B)^3 \end{aligned}$$

shows that

$$N_{U,H}(B) = \sum_{\boldsymbol{\eta} \in \mathcal{E}^*(B)} \frac{Y_1Y_5Y_6}{\eta_2} G_2(Y_0)\vartheta(\boldsymbol{\eta}) + O(B(\log B)^{4-1/5}).$$

Finally, we note that

$$\frac{Y_1Y_5Y_6}{\eta_2} G_2(Y_0) = \frac{Y_1Y_5Y_6}{\eta_2 Y_0^2} \omega_\infty = \frac{B}{\boldsymbol{\eta}^{(1,1,1,1)}} \omega_\infty$$

using Lemma 3. ☐

For $\mathbf{k} = (k_1, k_2, k_3, k_4) \in \mathbb{Z}^4_{>0}$, let

$$\Delta_{\mathbf{k}}(n) := \sum_{\boldsymbol{\eta} \in \mathbb{Z}^4_{>0}, \boldsymbol{\eta}^{(k_1,k_2,k_3,k_4)} = n} \frac{\vartheta(\boldsymbol{\eta})}{\boldsymbol{\eta}^{(1,1,1,1)}}.$$

Consider the Dirichlet series

$$F_{\mathbf{k}}(s) = \sum_{n=1}^{\infty} \frac{\Delta_{\mathbf{k}}(n)}{n^s} = \sum_{\boldsymbol{\eta} \in \mathbb{Z}^4_{>0}} \frac{\vartheta(\boldsymbol{\eta})}{\boldsymbol{\eta}^{(k_1s+1,k_2s+1,k_3s+1,k_4s+1)}}.$$

It is absolutely convergent for $\Re e(s) > 0$. We write it as an Euler product $F_{\mathbf{k}}(s) = \prod_p F_{\mathbf{k},p}(s)$, where we compute that $F_{\mathbf{k},p}(s)$ is

$$\begin{aligned} (1 - 1/p) \cdot \Bigg( &(1 + 1/p) + \frac{1 - 1/p}{p^{k_1s+1} - 1} + \frac{1 - 1/p}{p^{k_2s+1} - 1} + \frac{1 - 1/p}{p^{k_4s+1} - 1} \\ &+ \frac{1 - 1/p}{p^{k_3s+1} - 1} \left( (1 - 2/p) + \frac{1 - 1/p}{p^{k_1s+1} - 1} + \frac{1 - 1/p}{p^{k_2s+1} - 1} + \frac{1 - 1/p}{p^{k_4s+1} - 1} \right) \Bigg). \end{aligned}$$

For $\varepsilon > 0$ and $k \in \{(2,2,3,2),(3,3,4,2)\}$ and all $s \in \mathbb{C}$ lying in the half-plane $\Re e(s) \geqslant -1/8+\varepsilon$, we have

$$F_{\mathbf{k},p}(s) \prod_{j=1}^{4} \left(1 - \frac{1}{p^{k_j s+1}}\right) = 1 + O_\varepsilon(p^{-1-\varepsilon}).$$

We define

$$E_{\mathbf{k}}(s) := \prod_{j=1}^{4} \zeta(k_j s+1), \quad G_{\mathbf{k}}(s) := \frac{F_{\mathbf{k}}(s)}{E_{\mathbf{k}}(s)}$$

and note that $F_{\mathbf{k}}(s)$ has a meromorphic continuation to $\Re e(s) \geqslant -1/8+\varepsilon$ with a pole of order 4 at $s=0$.

As in [BD07, Lemma 15], we use a Tauberian theorem to show that

$$M_{\mathbf{k}}(t) := \sum_{n \leqslant t} \Delta_{\mathbf{k}}(n)$$

can be estimated as

$$\frac{G_{\mathbf{k}}(0)P(\log t)}{4!\prod_{j=1}^{4} k_j} + O(t^{-\delta})$$

for some $\delta > 0$ and $P$ a monic polynomial of degree 4.

Using Lemma 10 and the definitions of $\Delta_{\mathbf{k}}$ and $M_{\mathbf{k}}$,

$$\begin{aligned} N_{U,H}(B) &= \omega_\infty B \sum_{n \leqslant B} (\Delta_{(2,2,3,2)}(n) - \Delta_{3,3,4,2}(n)) + O(B(\log B)^{4-1/5}) \\ &= \omega_\infty G_{\mathbf{k}}(0) \frac{1}{4!}\left(\frac{1}{2^3 \cdot 3} - \frac{1}{2 \cdot 3^2 \cdot 4}\right) B(\log B)^4 + O(B(\log B)^{4-1/5}) \end{aligned}$$

Since

$$\alpha(\widetilde{S}) = \frac{1}{864} = \frac{1}{4!}\left(\frac{1}{2^3 \cdot 3} - \frac{1}{2 \cdot 3^2 \cdot 4}\right)$$

and

$$G_{\mathbf{k}}(0) = \prod_p \left(1 - \frac{1}{p}\right)^5 \left(1 + \frac{5}{p} + \frac{1}{p^2}\right),$$

this completes the proof of the theorem.

## References

[BB07] R. de la Bretèche and T. D. Browning, *On Manin's conjecture for singular del Pezzo surfaces of degree 4. I*, Michigan Math. J. **55** (2007), no. 1, 51–80. MR MR2320172

[BBD07] R. de la Bretèche, T. D. Browning, and U. Derenthal, *On Manin's conjecture for a certain singular cubic surface*, Ann. Sci. École Norm. Sup. (4) **40** (2007), no. 1, 1–50. MR MR2332351

[BD07] T. D. Browning and U. Derenthal, *Manin's conjecture for a quartic del Pezzo surface with* $\mathbf{A}_4$ *singularity*, arXiv:0710.1560, 2007.

[BT98] V. V. Batyrev and Yu. Tschinkel, *Manin's conjecture for toric varieties*, J. Algebraic Geom. **7** (1998), no. 1, 15–53. MR MR1620682 (2000c:11107)

[CLT02] A. Chambert-Loir and Yu. Tschinkel, *On the distribution of points of bounded height on equivariant compactifications of vector groups*, Invent. Math. **148** (2002), no. 2, 421–452. MR MR1906155 (2003d:11094)

[Der06] U. Derenthal, *Singular Del Pezzo surfaces whose universal torsors are hypersurfaces*, arXiv:math.AG/0604194, 2006.

- [Der07] U. Derenthal, *On a constant arising in Manin's conjecture for del Pezzo surfaces*, Math. Res. Lett. **14** (2007), no. 3, 481–489. MR MR2318651
- [DJT07] U. Derenthal, M. Joyce, and Z. Teitler, *A nef cone volume for generalized Del Pezzo surfaces*, arXiv:math.AG/0703202, 2007.
- [DT07] U. Derenthal and Yu. Tschinkel, *Universal torsors over del Pezzo surfaces and rational points*, Equidistribution in number theory, an introduction, NATO Sci. Ser. II Math. Phys. Chem., vol. 237, Springer, Dordrecht, 2007, pp. 169–196. MR MR2290499 (2007j:14024)
- [FMT89] J. Franke, Yu. I. Manin, and Yu. Tschinkel, *Rational points of bounded height on Fano varieties*, Invent. Math. **95** (1989), no. 2, 421–435. MR 89m:11060
- [Pey95] E. Peyre, *Hauteurs et mesures de Tamagawa sur les variétés de Fano*, Duke Math. J. **79** (1995), no. 1, 101–218. MR MR1340296 (96h:11062)

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT ZÜRICH, WINTERTHURERSTRASSE 190, 8057 ZÜRICH, SWITZERLAND
*E-mail address*: `ulrich.derenthal@math.unizh.ch`